



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

spilling and the reptile apparently thrived. Dissection after death showed blood-clots around the heart; otherwise all organs looked normal. Thick layers of fat lined the sides of the abdomen. A few days before, while feeding, the forceps slipped and the reptile's jaws closed with sufficient force to splinter the glass tube. Injury received through this accident probably caused death.

Shedding of the skin did not occur at regular intervals, but rather continuously, small patches peeling off here and there.

On October 26, 1907, a museum attendant was bitten in the fleshy part of the base of the thumb of the right hand. The reptile's hold was extremely tenacious, and some time, perhaps a minute, elapsed before the hand was released. The wound showed 6 small punctures. Swelling and discoloration of the hand, accompanied by great pain in the hand and arm, followed quickly. Although immediately placed under medical care, when the patient reported for duty two weeks later, he still complained of numbness in the hand and arm and occasional dizziness. Lack of mental reserve and a physical condition below the average in this case no doubt contributed in rendering the effect of the Gila Monster's bite unusually severe.

GEORGE P. ENGELHARDT,  
*Brooklyn, N. Y.*

## A HYBRID CENTRARCHID.

On December 12-14, 1911, a small collection of fishes was made in the Chesapeake and Ohio Canal above Violet's Lock, about 25 miles above the city of Washington, D. C.

Among the fishes taken was a Centrarchid which is apparently a cross between *Chænobryttus*

*gulosus* (Cuvier & Valenciennes) and *Lepomis gibbosus* (Linnæus). At the time of capture it was noted that the form and color were unusual and unlike any of the Centrarchids common to these waters.

Following is a table of comparative measurements of the hybrid and a typical example of each of the other species of nearly the same size :

	<i>Chænobryttus gulosus</i>	<i>Hybrid</i>	<i>Lepomis gibbosus</i>
Total length in cm.....	19.4	16.3	15.4
Head (without flap) in standard length.....	2.67	2.59	2.91
Depth in standard length.....	2.28	2.10	2.10
Eye in head.....	5.28	4.63	4.00
Snout in head.....	3.86	3.00	3.65
Maxillary.....	2.15	2.55	3.23
Maxillary reaching.....	to behind pupil	to front of pupil	to front of eye
Interorbital.....	4.06	3.64	3.11
Pectoral.....	1.66	1.42	1.20
Teeth on tongue.....	present	present	absent
Dorsal formula.....	X, 10, 1	X, 10, 1	X, 11, 1
Anal formula.....	III, 9, 1	III, 9, 1	III, 10, 1
Scale formula .....	7+11-42 (+5)	7+12-42 (+5)	7+12-39 (+4)
Rows of scales on cheeks .....	6	6	5

The profile of the hybrid is very different from *C. gulosus*, the body being short and deep, ovate; the snout is long, pointed, and the upper profile, from tip of snout to origin of dorsal, is relatively straight.

In *C. gulosus* the pectoral is broad, distal margin evenly rounded; in the hybrid, it is broad, pointed, upper rays longest, and in *L. gibbosus* it is narrow, pointed, median rays longest. Although the color pattern of the hybrid is nearest to that of *C. gulosus*, it is unlike that of any example seen in these waters, the color markings are less distinct

and on lower side and belly blend into those of typical examples of *L. gibbosus*. The coloration of the fins and the form and coloration of the opercular flap agree with *C. gulosus*.

The writer has seen other hybrid Centrarchids from the vicinity of Washington, D. C., one of these being apparently a cross between *C. gulosus* and *L. gibbosus*; another between *C. gulosus* and *L. cyanellus* Rafinesque. In this connection it is interesting to note that *C. gulosus* is not a native species, having been introduced into the Potomac by the Fish Commission about 1895.

LEWIS RADCLIFFE,  
*U. S. Bureau of Fisheries.*

## SOME NOTES ON THE CRICKET FROG ON LONG ISLAND.

The country lying between Flushing and Jamaica is an excellent place for frogs. In the woods and between the woods are many small ponds varying from a small pool to a fair-sized pond, big and deep enough to swim in. Here as early as March 23, 1913, I found and collected several Cricket Frogs. Later, as the spring advanced, they became quite common, the small gray frogs being the most plentiful.

This year, 1914, their appearance was later, April 19 being the first. Two weeks later, May 3, I collected several and heard the song; about 11 o'clock one sang as I watched it. Later in the day, at a pond some two miles northeast, I heard some more, while those in my collecting box sang several times.

HOWARTH S. BOYLE,  
*Elmhurst, N. Y.*